Digital Aerial Baseline Survey of Marine Wildlife

In Support of New York State Offshore Wind Energy

Greg Lampman NYSERDA
Greg Forcey Normandeau Associates, Inc.
Steph McGovern APEM

NORMANDEAU ASSOCIATES
Environmental Consultants
APEM

Outline

- Normandeau/APEM Experience
- Proposal Framework
- Known Wildlife Distributions in OPA
- Approach
  - Camera Sensor
  - Flight Planning and Survey Design
  - Data Output
  - Adaptive Methods Consideration
  - Survey Timing
- Data Distribution

Technology Evolution

- **Europe 2007:** Aerial digital surveys are used for collecting offshore biological data
- **USA 2011:** On behalf of BOEM, Normandeau completed a comparison of **three offshore survey methodologies**
  - Boat-based visual
  - Low-altitude aerial visual
  - High-altitude aerial digital

NORMANDEAU ASSOCIATES
Environmental Consultants
APEM

Survey Vessels

Boat surveys:
40-ft sport-fishing boat

Aerial surveys:
Cessna 337 Skymaster aircraft

The digital imagery camera plane flew between 450 and 1000 m altitude

NORMANDEAU ASSOCIATES
Environmental Consultants
APEM

Comparison-Density

- **Turtle Density Estimates**
  - Digital survey estimates 4x higher than visual aerial data
  - Digital survey estimates 10x higher than boat survey data

- **Reasons**
  - Low visibility of turtles from boats at sea-level and from aircraft given the short observation time available
  - Disturbance by both boat and aerial visual survey platforms

Comparison-Identification

- **Birds:** digital aerial surveys and boat-based surveys achieved higher success than visual aerial surveys
- **Turtles and Cetaceans:** boat-based surveys had highest success



NORMANDEAU ASSOCIATES
Environmental Consultants
APEM

Conclusions

- Digital aerial surveys offer a significant methodological improvement over visual surveys for turtles
- For marine birds and mammals in low-density situations, digital and observer-based survey methods may produce similar density estimates
  - *Caveat:* No marine mammals with long dive times (e.g., whales) were recorded in this study
- Density calculations from digital aerial surveys are more accurate and precise
  - Precise definition of areal coverage
  - Reduced animal repulsion/attraction effects
  - Not affected by observer detectability biases

APEM Experience

- Provided aerial surveys for 10 years
  - 40 staff in remote sensing team
  - Own aircraft & camera systems
- Extensive experience in nearshore and offshore habitats
- Examples of areas worked in Europe:
  - North Sea
  - German N Sea
  - Irish Sea
  - Baltic Sea
  - Pentland Firth and Orkney Waters
  - Carmarthen Bay

NORMANDEAU ASSOCIATES
Environmental Consultants
APEM

Innovations & Collaboration

- Innovative approach to survey designs
  - Camera system allows flexible approach
  - E.g., development and application of quasi-random survey to estimate seabird avoidance to operational wind turbines
- APEM-Normandeau ornithology and marine mammal collaborations
  - Baryonyx, Gulf of Mexico
  - Confidential Project (onshore breeding bird census)

NORMANDEAU ASSOCIATES
Environmental Consultants
APEM

Known Wildlife Distributions in OPA

NORMANDEAU ASSOCIATES
Environmental Consultants
APEM

Data Sources: Spatial Data

- Northwest Atlantic Seabird Catalog by USGS, BOEM, available at NOAA
- New England Aquarium using the North Atlantic Right Whale Consortium
- MARCO Mid-Atlantic Ocean Data Portal

Spatial: Birds



NY Offshore Planning Area

NY Wind Energy Area

Atlantic Offshore Seabird Dataset Catalog

- Atlantic Puffin
- Razorbill
- Audubon's Shearwater
- Cory's Shearwater
- Red-throated Loon
- Black-legged Kittiwake
- Northern Gannet
- Herring Gull
- Jaegers
- Scoters

NORMANDEAU ASSOCIATES Environmental Consultants

APEM

Spatial: Roseate Terns



NORMANDEAU ASSOCIATES
Environmental Consultants
APEM

Spatial: Turtles





NORMANDEAU ASSOCIATES
Environmental Consultants
APEM

Approach

NORMANDEAU ASSOCIATES
Environmental Consultants
APEM

Innovations & Collaboration

- Innovative approach to survey designs
  - Camera system allows flexible approach
  - E.g., development and application of quasi-random survey to estimate seabird avoidance to operational wind turbines
- APEM-Normandeau ornithology and marine mammal collaborations
  - Baryonyx, Gulf of Mexico
  - Confidential Project (onshore breeding bird census)

NORMANDEAU ASSOCIATES
Environmental Consultants
APEM

Camera Sensor

- Shearwater III
- 1.5 cm for NYSERDA
- Integrated GPS and IMU

Key Advantages to 1.5 cm Resolution

- 1.5 cm resolution is the same as a nickel at sea-surface
- High quality
- More accurate measurements
  - Head to tail and wingspan
  - Flight height calculation
- Greater confidence in ID
  - Measurements (e.g., flying shearwaters, sitting auks, and gulls)
  - Tern bill colour visible
  - Increased chance of ID'ing individual mammals

Example Bird ID Rates

2 cm v. predicted 1.5 cm

Species Group	Species	% ID (2 cm)	% ID (1.5 cm)
Ducks	Black Scoter	>85%	>98%
	Red-breasted Merganser	>80%	>98%
	Long-tailed Duck	>85%	>95%
Loons	Common & Red-throated Loons	>90%	>98%
Petrels and Shearwaters	Wilson's Storm Petrel	>75%	>85%
	Great & Cory's Shearwater	>80%	>90%
	Manx, Audubon's & Sooty Shearwaters	>60%	>80%
Jeagers and Skuas	Parasitic & Pomarine Jaeger	>85%	>95%
Gulls	Lesser & Great Black-backed Gulls	100%	100%
	Herring Gull	>90%	>99%
	Kittiwake, Laughing & Bonaparte's Gull	>85%	>95%
Terns	Caspian & Royal Terns	>85%	>95%
	Common, Roseate & Forster's Terns	>60%	>75%
Auks	Common Murre & Razorbill	>90% Summer, <50% in Winter	>99% in Summer, <50% in Winter
	Dovekie & Atlantic Puffin	>60%	>90%

NORMANDEAU ASSOCIATES Environmental Consultants APEM

Example High-Res Bird Snags

NORMANDEAU ASSOCIATES
Environmental Consultants

APEM

Flight Planning

- Twin engine capable of flying less than 140 knots
- Proposed to fly at around 1,500 ft
  - Safety
  - Weather
  - Resolution
  - Coverage

Survey Design
Confidence in Estimates

- Without small confidence intervals, it is impossible to assess if environmental change has had a significant impact on a population
  - High confidence = small confidence intervals
  - Low confidence = large confidence intervals





NORMANDEAU ASSOCIATES
Environmental Consultants
APEM

Survey Design

- OPA = 43,650 km^2
- 7% Transect of OPA (3,073 km^2)

  584 m x 110 m ground sampling per image

- 10% Grid of WEA + 4 km buffer (128 km^2)

  330 m x 219 m ground sampling per image

- Rationale for Transect and Grid Design

Survey Design









NORMANDEAU ASSOCIATES
Environmental Consultants
APEM

Survey Design



NORMANDEAU ASSOCIATES
Environmental Consultants
APEM

Survey Design
Grid versus Transect

- Both designs have benefits and drawbacks
- Key features transect design
  - Suitable for providing a generalisation of habitat and species abundance over large areas
  - Quicker to collect data
- Key features of grid design
  - Power to detect and monitor change in abundance
  - Greater number of replicates
  - Uniform sampling across environmental gradients

Survey Design
Accommodating Glare

- **Consideration will be given to**
  - Sun angle (time of day / time of year)
  - Sea state (4 or less aiming for 2 or less)
  - Direction of flight in relation to the sun
  - Camera angle
  - Camera technicians continuously monitor the images collected for quality. Image acquisition will be stopped until suitable conditions occur
  - Extra imagery is routinely collected to replace a few expected glint-affected images

Data Output
Geospatial Accuracy

- Custom flight planning software pre-programs the survey transects and grids
- System-specific, aircraft mounted GPS/GNSS systems ensure that surveys are flown as accurately as possible
- Automatic image acquisition over specified locations
- As data capture occurs, GPS data are automatically logged with each exposure including the xyz coordinate and heading of the camera at the point of capture



Data Output

- Identify wildlife to lowest possible taxonomic group
  - Birds
  - Marine mammals
  - Turtles
  - Sharks
  - Rays
  - Fish
- Geo-rectified image snags with associated metadata including height and direction
- Other Anthropogenic Data
  - Boats, met masts, etc.

Data Output

Flight Height





NORMANDEAU ASSOCIATES Environmental Consultants

APEM

Data Output
Flight/Path Direction

- Individual wildlife are geo-referenced
- Bearing is automatically determined from head-tail axis
- Extraction to GIS
- Rose diagrams produced for defined areas



- Predominant flight/path direction is detectable

NORMANDEAU ASSOCIATES
Environmental Consultants
APEM

Data Output
Flight/Path Direction



Adaptive Methods Consideration

- Goal→ Create best value for 3 year data set
- Do Year 1 and/or 2 data confirm historical hot spot and cold spots?
- Stratified by area and/or by target species
- More detailed data–sampling emphasis and/or resolution

Data Sources: Temporal Data

- Northwest Atlantic Seabird Catalog by USGS, BOEM, available at NOAA
- New England Aquarium using the North Atlantic Right Whale Consortium
- MARCO Mid-Atlantic Ocean Data Portal
- eBird

Temporal: Marine Mammals



MAR_APR JUL OCT_NOV

NORMANDEAU ASSOCIATES Environmental Consultants APEM

Temporal: Marine Mammals



MAR_APR JUL OCT_NOV

NORMANDEAU ASSOCIATES Environmental Consultants APEM

Temporal: Turtles



NORMANDEAU ASSOCIATES
Environmental Consultants
APEM

Temporal: eBird



NORMANDEAU ASSOCIATES
Environmental Consultants
APEM

Sensitive Species

- **Population Sensitivity**
  - Roseate Tern
  - Cory's Shearwater
  - Audubon's Shearwater
- **Displacement Sensitivity**
  - Roseate Tern
  - Atlantic Puffin
  - Razorbill
  - Red-throated Loon
  - Northern Gannet
- **Collision Sensitivity**
  - Roseate Tern
  - Herring Gull
  - Jaegers
  - Black-legged Kittiwake

Robinson Willmott, J. C., G. Forcey, and A. Kent. 2013. *The relative vulnerability of migratory bird species to offshore wind energy projects on the Atlantic Outer Continental Shelf*. OCS Study BOEM 2013-207.

Survey Timings: Winter

February-March







NORMANDEAU ASSOCIATES
Environmental Consultants
APEM

Survey Timings: Winter

February-March



NORMANDEAU ASSOCIATES
Environmental Consultants
APEM

Survey Timings: Spring
April-May



NORMANDEAU ASSOCIATES
Environmental Consultants
APEM

Survey Timings: Spring

April-May



NORMANDEAU ASSOCIATES
Environmental Consultants
APEM

Survey Timings: Summer

July-August





NORMANDEAU ASSOCIATES
Environmental Consultants
APEM

Survey Timings: Summer

July-August



NORMANDEAU ASSOCIATES
Environmental Consultants
APEM

Survey Timings: Fall
October-November









NORMANDEAU ASSOCIATES
Environmental Consultants
APEM

Survey Timings: Fall

October-November



NORMANDEAU ASSOCIATES
Environmental Consultants
APEM

Survey Timing
Proposed

Seasonal Survey	Months
Winter	February/March
Spring	End of April–May
Summer	End of July–August
Fall	October/November

Data Distribution

AMAPPS program NOAA and USFWS

Northwest Atlantic Seabird Catalog by USGS, BOEM, available at NOAA

North Atlantic Right Whale Consortium

MARCO Mid-Atlantic Data Portal

MA Marine Fisheries

NY DOS Geographic Information Gateway

Large Pelagics Research Center at UMass

Feedback and Suggestions

Feedback and suggestions should be sent to:

Gregory.lampman@nyserda.ny.gov

http://remote.normandeau.com/NYSERDA

NORMANDEAU ASSOCIATES
Environmental Consultants
APEM

WREN Webinars Held to Date

- This is WREN's 8th webinar.
- Each offers perspectives from multiple nations on research, monitoring, methodologies, and results of wind energy and wildlife interactions.
- If you are not on the webinar mailing list, simply send a blank email to **join-wind-webinars@lyris.pnnl.gov**

All webinars and associated presentations are archived on the WREN Hub at: http://tethys.pnnl.gov.

Thank You!

For questions or to provide ideas for future webinars, please contact one of the members of the WREN Operating Agent team:

- **Karin Sinclair karin.sinclair@nrel.gov**
- **Andrea Copping andrea.copping@pnnl.gov**
- **Jocelyn Brown-Saracino jocelyn.brown-saracino@ee.doe.gov**

iea wind